

BIOGRAPHICAL SKETCH

Provide the following information for the key personnel in the order listed for Form Page 2.
Follow the sample format for each person. **DO NOT EXCEED FOUR PAGES.**

NAME		POSITION TITLE		
GATTI, RICHARD, A.		PROFESSOR STEP IX		
EDUCATION/TRAINING (Begin with baccalaureate or other initial professional education, such as nursing, and include postdoctoral training)				
INSTITUTION AND LOCATION		DEGREE (if applicable)	YEAR (s)	FIELD OF STUDY
Columbia College, New York, New York		B.A.	1958	Pre-Med/Music
St. Louis U., School of Medicine, St. Louis		M.D.	1962	Medicine
Children's Memorial Hospital, Northwestern, Chicago			1962-1965	Pediatrics
U. of Minnesota, Minneapolis, MI			1968-1972	Immunology
Karolinska Inst., Tumor Biology, Stockholm			1972-1974	Tumor Biology
			1965-1966	

RESEARCH AND PROFESSIONAL EXPERIENCE:

USPHS Traineeship in Biochemical Genetics (Northwestern), U.S. Army Medical Corps, Captain (Eritrea, Ethiopia)	[David Y-Y. Hsia]	1965-1966
USPHS Traineeship in Immunogenetics (U. of Minnesota)	[Robert A. Good]	1966-1968
USPHS Special Fellowship (U. of Minnesota)	[Robert A. Good]	1968-1969
USPHS Research Career Development Award (Minn./Sweden)	[Robert A. Good/Georg Klein]	1969-1971
Cedars-Sinai Medical Center, Los Angeles Director, Pediatric Hematology, Oncology & Immunology		1974-1980
UCLA School of Medicine, Los Angeles Professor, Department of Pediatrics		1975-1980
Professor, Department of Pathology & Laboratory Medicine		1980-
Co-director, Diagnostic Molecular Pathology		1986-

BIBLIOGRAPHY (selected from over 300 references)

- Gatti** RA, et al. Hereditary lymphopenic agammaglobulinemia associated with a distinctive form of short-limbed dwarfism and ectodermal dysplasia. *J Pediat* 75:675-684,1969.
- Gatti** RA, Meuwissen HJ, Allen HD, Hong R, Good RA: Immunologic reconstitution of sex-linked lymphopenic immunologic deficiency by allogeneic bone marrow transplantation. *Lancet* II:1366, 1968.
- Garrioch DB, Good RA, **Gatti** RA: Lymphocyte response to PHA in patients with nonlymphoid tumors. *Lancet* I:618, 1970.
- Gatti** RA, Good RA: The immunological deficiency diseases. *Medical Clinics of North Amer.* Ed. MG Wohl, WB Saunders, Pub. Vol. 54, 1970, pp 281-307.
- Gatti** RA, Good RA: Occurrence of malignancy in immunodeficiency diseases. *Canc.* 28:89,1971.
- Gatti** RA, Meuwissen HJ, Terasaki PI, Good RA: Recombination within the HLA locus. *Tissue Antigens* 1:239-241, 1971.
- Gatti** RA: On the classification of patients with primary immunodeficiency disorders. *Clin Immunol Immunopath* 3:243, 1974.
- Walford RL, Gossett T, Troup GM, **Gatti** RA, Mittal KA, Robbins A, Ferrar GB, Zeller E: The Meritt alloantigenic system of human B-lymphocytes: Evidence of thirteen possible factors including one six-member segregant series. *J Immunol* 116:1704, 1976.
- Gatti** RA, et al.: Ataxia-telangiectasia: A multiparameter analysis of eight families. *Clin Immunol Immunopath* 23:501-516, 1982.
- Gatti** RA, Concannon P, Salser W: Multiple use of Southern blots. *BioTechniq.* 2:148-155, 1984.
- Vinters HV, **Gatti** RA, Rakic P: Sequence of cellular events in cerebellar ontogeny relevant to expression of neuronal abnormalities in ataxia-telangiectasia. In: Ataxia-telangiectasia: Genetics, Neuropathology, and Immunology of a Degenerative Disease of Childhood. Eds. RA **Gatti** and M Swift. Alan R Liss, Inc., New York. 1985. Pp. 233-255.
- Concannon P, **Gatti** RA, Hood LE: Human TCR V-beta gene polymorphism. *J Exp Med* 165:1130-1140, 1987.
- Gatti** RA, Shaked R, Mohandas TK, Salser W: Human ferritin genes: chromosomal assignments and polymorphisms. *Am J Hum Genet* 41:654-667, 1987.
- Gatti** RA, et al: Localization of an ataxia-telangiectasia gene to chromosome 11q22-23. *Nature* 336, 577-580, 1988.
- Jaspers MGJ, **Gatti** RA, Baan C, Bootsma D: Genetic complementation analysis of ataxia-telangiectasia and Nijmegen breakage syndrome: a survey of 50 patients. *Cytogenet Cell Genet* 49:259-263,1988.
- Curry CJR, et al: ATFresno: a phenotype linking ataxia-telangiectasia with Nijmegen breakage syndrome. *Am J Hum Genet* 45:270-275, 1989.
- Kojis TL, Schreck RR, **Gatti** RA, Sparkes RS: Tissue specificity of chromosomal rearrangements in ataxia-telangiectasia. *Human Genetics* 83:347-352, 1989.
- Charmley P, et al: A primary linkage map of the human chromosome 11q22-23 region. *Genomics* 6:316-323, 1990.
- Wei S, Rocchi M, Archidiacono N, Sacchi N, Romeo G, **Gatti** RA: Physical mapping of the human chromosome 11q23 region containing the ataxia-telangiectasia locus. *Cancer Genetics Cytogenet* 46: 1-8, 1990.

Charmley PR, Chao A, Concannon P, Hood L, **Gatti RA**: Haplotyping the gene complex of the human T-cell receptor beta-chain using DNA polymorphisms. *Proc. Natl. Acad. Sci. USA.* 87: 4823-4827, 1990.

Sanal O, et al: Further mapping of an ataxia-telangiectasia locus to the chromosome 11q23 region. *Am J Hum Genet* 47: 860-866, 1990.

Gatti RA, Boder E, Vinters HV, Sparkes RS, Norman A, Lange K: Ataxia-telangiectasia: an interdisciplinary approach to pathogenesis. *Medicine* 70: 99-117, 1991.

Foroud T, et al: Localization of an ataxia-telangiectasia locus to a 3-cM interval on chromosome 11q23: linkage analyses of 111 families by an international consortium. *Amer J Hum Genet* 49:1263-1279, 1991.

Gatti RA, et al: Prenatal genotyping of ataxia-telangiectasia. *Lancet* 342: 376, 1993.

Gatti RA and Painter RB, Eds. Ataxia-Telangiectasia. Springer-Verlag, Heidelberg. 1993 Pp 1-283.

Uhrhammer, N, Concannon P, Huo Y, Nakamura Y, **Gatti RA**. A pulsed-field gel electrophoresis map in the A-T region of chromosome 11q22.3. *Genomics* 20:278-280, 1994.

Huo YK, Wang Z, Hong J-H, Chessa L, McBride WH, Perlman SL, **Gatti RA**. Radiosensitivity of ataxia-telangiectasia, X-linked agammaglobulinemia and related syndromes. *Canc Res.* 54: 2544-2547, 1994

Gatti RA, McConville CM, Taylor AMR. Sixth International Workshop on A-T. *Canc Res* 54: 6007-6010, 1994.

Lange E, et al. Localization of an ataxia-telangiectasia gene to a ~500 kb interval on chromosome 11q23.1: linkage analysis of 176 families in an international consortium. *Amer J Hum Genet* 57:112-119, 1995..

Uhrhammer N, et al. Sublocalization of an ataxia-telangiectasia gene distal to D11S384 by ancestral haplotyping in Costa Rican families. *Amer J Hum Genet* 57:103-111, 1995.

Savitsky et al. A single ataxia telangiectasia gene with a product similar to PI-3 kinase. *Science* 268: 1749-1753, 1995.

Stumm M, **Gatti RA**, Reis A, Chrzanoska K, Seemanova E, Sperling K, Wegner R-D. The ataxia-telangiectasia-variant genes 1 and 2 are distinct from the AT gene on chromosome 11q23.1 *Amer J Hum Genetics* 57:960-962, 1995.

Telatar M, Wang Z, Udari N, Liang T, Concannon P, Lavin M, Shiloh Y, Good RA, **Gatti RA**. Ataxia-telangiectasia: mutations in ATM cDNA detected by protein truncation screening. *Amer J Hum Genet* 59:40-44, 1996.

Wright J, Teraoka S, Onengut S, Tolun A, **Gatti RA**, Ochs HD, Concannon P. A high frequency of distinct ATM gene mutations in ataxia-telangiectasia. *Amer J Hum Genet* 59:839-846, 1996.

Chen X, et al. CAND3: a ubiquitously-expressed gene immediately adjacent and in opposite transcriptional orientation to the ATM gene at 11q23.1 *Mamm Genome.* 8:129-133,1997.

Concannon P, **Gatti RA**. Diversity of ATM gene mutations detected in patients with ataxia-telangiectasia. *Human Mutation* 10:100-107,1997.

Telatar M et al. Ataxia-telangiectasia: identification and detection of founder-effect mutations in the ATM gene in ethnic populations. *Am J Hum Genet* 62: 86-97, 1998.

Cerosaletti KM, et al. Fine localization of the Nijmegen Breakage Syndrome gene to 8q21: evidence for a common founder haplotype. *Am J Hum Geneet* 63: 125-134, 1998.

Varon R, et al. Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen Breakage Syndrome. *Cell* 93:467-476,1998.

Castellvi-Bel S. et al. New mutations, polymorphisms, and rare variants in the ATM gene detected by a novel SSCP strategy. *Human Mutation* 14:156-162,1999.

Teraoka SN, et al. Splicing defects in the ataxia-telangiectasia gene, ATM: underlying mutaitons and consequences. *Am J Hum Genet* 64:1617-1631,1999.

Lavin MF, Concannon P, **Gatti RA**. Eighth International Workshop on Ataxia-Telangiectasia (ATW8). *Canc Res.* 59:3845-3849,1999.

Gatti RA, Tward A, Concannon P. Cancer risk in ATM heterozygotes: a model of phenotypic and mechanistic differences between missense and truncating mutations. *Molec Biol Metab.* 68: 419-423,1999.

Gatei M et al. ATM-dependent phosphorylation of nibrin in response to radiation exposure. *Nature Genet* 25:115-119, 2000.

Zhao S, et al. A functional link between ATM kinase and NBS1 in the DNA damage response. *Nature* 405:473-476,2000

Becker-Catania SG, et al. Ataxia-Telangiectasia: phenotype/genotype studies of ATM protein expression, mutations, and radiosensitivity. *Molec Genetics Metabolism.* 70: 122-133, 2000.

Hiel JA, et al (International NBS Study Group). Nijmegen Breakage Syndrome. *Arch Dis Child* 82:400-406, 2000.

Buzin CH et al. Scanning by DOVAM-S detects all unique sequence changes in blinded analyses: evidence that the scanning conditions are generic. *BioTechniques* 28:746-750, 2000.

Yeo TC, Xia D, Hassouneh S, Yang XO, Sabath DE, Sperling K, **Gatti RA**, Concannon PJ, Willerford DM. V(D)J rearrangement in Nijmegen Breakage Syndrome. *Mol Immunol* 37:1131-1139, 2000.

Gatti RA. The Inherited Basis of Human Radiosensitivity. *Proc. of Nobel Foundation Forum. Acta Oncologica.* 40:702-711, 2001.

O'Driscoll M et al. DNA ligase IV mutations identified in patients exhibiting developal delay and immunodeficiency. *Molecular Cell.* 8: 1175-1185, 2001.

Chun HH et al. TCL-1, MTCP-1 and TML-1 gene expression profile in ataxia-telangiectasia patients with clonal proliferations or leukemia. *International J Cancer.* 97: 726-731, 2002.

Sun X et al. Early diagnosis of ataxia-telangiectasia using radiosensitivity testing. *J Pediat* 40:724-731, 2002.

Gatti RA. Ataxia-Telangiectasia. In: The Metabolic and Molecular Bases of Inherited Disease. Edition 8. Scriver CR, Beaudet AL, Sly WS and Valle D. Publ. McGraw-Hill. 2001. New York. Pp. 705-732.

Pan-Hammarstrom Q, et al. ATM is not required in somatic hypermutation of VH, but is involved in the introduction of mutations in the switch mu region. *J. Immunol.* 170:3707-3716, 2003

Mitui, M et al. Independent mutational events are rare in the ATM gene: haplotype prescreening enhances mutation detection rate. *Hum Mutation* 22:43-50, 2003

Eng L, Coutinho G, Nahas S, Yeo G, Tanouye R, Babaei M, Dork T, Burge C, Gatti RA. Non-classical splicing mutations in the coding and non-coding regions of the ATM gene: maximum entropy estimates of splice junction strengths. *Hum Mutation* 23: 67-76, 2004.

Lai C-H et al: Correction of ATM gene function by aminoglycoside-induced readthrough of premature termination codons. *PNAS*, in press